arXiv:1107.1424v1 [gr-qc] 7 Jul 2011

Quantum corrections to gravity

Yukio Tomozawa
Michigan Center for Theoretical Physics, Randall Laboratory of Physics,
University of Michigan, Ann Arbor, MI. 48109-1040

(Dated: July 8, 2011)

This paper revisits quantum corrections to gravity. It was shown previously by other authors that quantum field theories in curved space time provide quadratic curvature forms as quantum corrections to gravity in a conformally flat metric. Application to a spherically symmetric and static (SSS) metric shows that only the Gauss Bonnet combination (GB) yields the correct expression. Using a variational method, the author shows that the metric he obtained in 1985 as an example in a simplified case was indeed the exact solution for a SSS metric. This proves that gravity becomes repulsive at short distances by quantum corrections.

PACS numbers: 04.70.-s, 95.85.Pw, 95.85.Ry, 98.54.Cm

I. INTRODUCTION

Quantum field thories on a curved space time were extensively studied in 1970-80 in the case of a conformally flat (CF) metric, and it was concluded that quantum corrections to gravity are expressed as quadratic curvature forms in that metric[1]. The application of that formalism to the Friedman Walker (FW) metric gave the interesting observation that a collapsed universe will bouce back to an expansion in some cases[2]. This provides a hint that quantum corrections to gravity might yield a repulsive component at short distances. The author studied quantum corrections for a SSS metric and obtained a repulsive force at short distance in the special case of a Gauss Bonnet (GB) combination, and also by a numerical computation. Based on this analysis, the author proposed a model for high energy cosmic rays produced from AGN in 1985[3], [4], [5]. This turns out to predict recent data from the Pierre Auger Project that suggests a possible correlation between high energy cosmic rays and locations of AGN[6]. Further discussion of the model yielded a new mass scale at the knee energy of the cosmic ray spectrum and a prediction for the mass of a dark matter particle, which is supported by recent data from HESS[7], [8], [9]. Because of successful predictions that agree with observational reality, it is important to solidify a theoretical basis for the model. From this point of view, the author has revisited the subject of quantum corrections on gravity. Using an explicit form for a SSS metric, it is shown that among proposed expressions in a conformally flat metric, only the Gauss Bonnet combination satisfies the finiteness of renormalization. This implies that the metric obtained by the author in 1985 is in fact the rigorous solution for quantum corrections to gravity and the repulsive nature of gravity at short distances has been established. A derivation of the final result by a variational method is presented.

II. QUANTUM CORRECTIONS IN A SSS METRIC

Extensive studies of quantum corrections to gravity in a CF metric can be summarized [1], [5] as

$$H^{\mu\nu} = \frac{1}{(n-4)}\frac{1}{\sqrt{|g|}}\frac{\delta}{\delta g_{\mu\nu}}\int F\sqrt{|g|}d^n x, \qquad (1)$$

where

$$F = AR^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta} + BR^{\alpha\beta}R_{\alpha\beta} + CR^2 \qquad (2)$$

$$= f_0 + (n-4)f_1 + O((n-4)^2), \qquad (3)$$

and A, B and C are constants. It is well known that the GB combination

$$A = 1, B = -4, C = 1 \qquad (4)$$

is finite for any metric. For a CF metric, the combinations

$$(A, B, C) = (0, 1, 0) \qquad (5)$$

and

$$(A, B, C) = (0, 0, 1/3) \qquad (6)$$

give finite values and contribute to a quantum correction term, although the two yield an identical correction. Using an explicit expression for curvature in n dimensions, I will examine this property for a SSS metric in n dimensions

$$ds^2 = e^{\nu}dt^2 - e^{\lambda}dr^2 - r^2 d\Omega^2, \qquad (7)$$

where ν and λ are functions of the radial coordinate, r, and the angular part can be expressed symbolically as

$$d\Omega^2 = d\omega_1^2 + d\omega_2^2 + \cdots + d\omega_{n-2}^2. \qquad (8)$$

From the computation of curvature in the appendix, one can express the integrand in Eq. (3), where f_0 and

f_1 are coefficients in the expansion in (n-4) and

$$\begin{aligned} f_0 &= (A+B/2+C)g_1 + (B+4C)g_2 \\ &\quad + (\nu'^2+\lambda'^2)(e^{-\lambda}/r)^2(2A+3B/2+4C) \\ &\quad - 4Cg_3. \end{aligned} \tag{9}$$

where

$$g_1 = (\nu'' + \nu'^2/2 - \nu'\lambda'/2)^2 e^{-2\lambda} + 4((e^{\lambda}-1)e^{-\lambda}/r^2)^2, \tag{10}$$

$$\begin{aligned} g_2 &= ((\nu'' + \nu'^2/2 - \nu'\lambda'/2)(\nu'-\lambda') - \nu'\lambda'/r \\ &\quad - 2(\nu'-\lambda')(e^{\lambda}-1)/r^2)e^{-2\lambda}/r, . \end{aligned} \tag{11}$$

and

$$g_3 = ((\nu'' + \nu'^2/2 - \nu'\lambda'/2)(e^{\lambda}-1) + \nu'\lambda')e^{-2\lambda}/r^2. \tag{12}$$

First manipulate the last term. Multiplying by the r dependent term,

$$\sqrt{|g|} = e^{(\nu+\lambda)/2}r^{n-2} \tag{13}$$

and noticing that a total derivative does not contribute to a variation, one can reformulate

$$\begin{aligned} G &= (\nu'' + \nu'^2/2)(e^{\lambda}-1)e^{-2\lambda}e^{(\nu+\lambda)/2}r^{n-4} \\ &= 2(e^{\nu/2})''(e^{-\lambda/2} - e^{-3\lambda/2})r^{n-4} \\ &= (2(e^{\nu/2})'(e^{-\lambda/2} - e^{-3\lambda/2})r^{n-4})' \\ &\quad + \nu'\lambda'(e^{\lambda}-3)/2e^{-2\lambda}e^{(\nu+\lambda)/2}r^{n-4} \\ &\quad - (n-4)\nu'(e^{\lambda}-1)e^{-2\lambda}e^{(\nu+\lambda)/2}r^{n-5} \end{aligned} \tag{14}$$

then

$$\begin{aligned} & - 4Cg_3e^{(\nu+\lambda)/2}r^{n-2} \\ &= 4C(n-4)\nu'(e^{\lambda}-1)e^{-2\lambda}e^{(\nu+\lambda)/2}r^{n-5}, \end{aligned} \tag{15}$$

where all terms cancel or a noncontributing total derivative term is dropped.

Now coming back to the expression for F, the first two terms should vanish independently, since they are the fourth derivative and third derivative of the function, ν. The condition for vanishing of the two coefficients

$$A + B/2 + C = 0 \tag{16}$$

and

$$B + 4C = 0 \tag{17}$$

yields a multiple of the GB condition, Eq. (4). For that, the third term automatically vanishes

$$\begin{aligned} & 2A + 3B/2 + 4C \\ &= 2(A+B/2+C) + (B+4C)/2 = 0 \end{aligned} \tag{18}$$

The fourth term is proportional to $(n-4)$ as is seen in Eq. (15). This term should be added to the quantum correction term f_1 in the variational calculation. This proves that quantum corrections for a SSS metric should consist of a GB combination exclusively, where

$$f_0 = 0. \tag{19}$$

Using expressions from the appendix, the expansion of F in $(n-4)$ [yield] yields

$$\begin{aligned} f_1 &= A((\nu'^2+\lambda'^2)(e^{-\lambda}/r)^2 + 6(e^{\lambda}-1)^2(e^{-\lambda}/r)^2) \\ &\quad + B(\frac{1}{2}(\nu'' + \nu'^2/2 - \nu'\lambda'/2)(\nu'-\lambda')e^{-2\lambda}/r \\ &\quad + \frac{5}{4}(\nu'^2+\lambda'^2)(e^{-\lambda}/r)^2 - \nu'\lambda'(e^{-\lambda}/r)^2/2 \\ &\quad - 3(\nu'-\lambda')((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2 \\ &\quad + 5((e^{\lambda}-1)/r)^2(e^{-\lambda}/r)^2) \\ &\quad + C(2(\nu'' + \nu'^2/2 - \nu'\lambda'/2)(\nu'-\lambda')e^{-2\lambda}/r \\ &\quad + 4(\nu'^2+\lambda'^2)(e^{-\lambda}/r)^2 - 8\nu'\lambda'(e^{-\lambda}/r)^2 \\ &\quad - 6(\nu'' + \nu'^2/2 - \nu'\lambda'/2)(e^{\lambda}-1)(e^{-\lambda}/r)^2 \\ &\quad - 16(\nu'-\lambda')((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2 \\ &\quad + 12(e^{\lambda}-1)^2(e^{-\lambda}/r^2)^2) \\ &\quad + 4C\nu'((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2, \end{aligned} \tag{20}$$

where the last term is added from Eq. (15). Rearranging appropriate terms, one gets

$$\begin{aligned} f_1 &= (\nu'^2+\lambda'^2)(e^{-\lambda}/r)^2(A + B/4 + B + 4C) \\ &\quad + (B+4C)g_4 \\ &\quad - 6Cg_3 \\ &\quad + ((e^{\lambda}-1)(e^{-\lambda}/r^2))^2(6A + 5B + 12C) \\ &\quad - (\nu'-\lambda')((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2(3B+16C) \\ &\quad + 4C\nu'((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2, \end{aligned} \tag{21}$$

where

$$g_4 = \frac{1}{2}((\nu'' + \nu'^2/2 - \nu'\lambda'/2)(\nu'-\lambda')r - \nu'\lambda')(e^{-\lambda}/r)^2 \tag{22}$$

In this expression, the first two terms vanish by the BG condition, Eq. (4), while the third term becomes higher order [of] in $(n-4)$ by Eq. (15). In the last two terms, the ν' terms cancel and finally one gets

$$f_1 = 4\lambda'((e^{\lambda}-1)/r)(e^{-\lambda}/r)^2 - 2(e^{\lambda}-1)^2(e^{-\lambda}/r^2)^2. \tag{23}$$

Multiplying by $e^{(\nu+\lambda)/2}r^2$, variation of

$$g_{00} = e^{\nu} \tag{24}$$

yields

$$H^{00} = \frac{1}{e^{(\nu+\lambda)/2}r^2}\frac{\delta}{\delta(e^{\nu})}(f_1e^{(\nu+\lambda)/2}r^2) \tag{25}$$

$$= \frac{1}{2}e^{-\nu}f_1 \tag{26}$$

or

$$\begin{aligned} H^0_0 &= \frac{1}{2} f_1 \qquad (27)\\ &= 2\lambda'((e^\lambda - 1)/r)(e^{-\lambda}/r)^2 - (e^\lambda - 1)^2(e^{-\lambda}/r^2)^2. \qquad (28)\end{aligned}$$

For the variation of

$$g_{11} = -e^\lambda, \qquad (29)$$

$$\begin{aligned} H^{11} &= \frac{1}{e^{(\nu+\lambda)/2} r^2}\Big(\frac{\delta}{\delta(-e^\lambda)}(f_1 e^{(\nu+\lambda)/2} r^2)\\ &- \big(\frac{\delta}{\delta(-(e^\lambda)')}(f_1 e^{(\nu+\lambda)/2} r^2)\big)'\Big)\\ &= \frac{1}{e^{(\nu+\lambda)/2} r^2}\Big(4(e^\lambda)' \frac{\frac{3}{2}e^{-5\lambda/2} - \frac{5}{2}e^{-7\lambda/2}}{r} e^{\nu/2}\\ &+ (e^{-\lambda/2} + 2e^{-3\lambda/2} - 3e^{-5\lambda/2})e^{\nu/2}\\ &+ 4\big(\frac{e^{-3\lambda/2} - e^{-5\lambda/2}}{r}\big)e^{\nu/2}\big)'\Big)\\ &= \frac{1}{e^{(\nu+\lambda)/2} r^2}\Big(2\nu' \frac{e^\lambda - 1}{r} e^{-3\lambda} e^{(\nu+\lambda)/2}\\ &+ (e^\lambda - 1)^2 e^{-3\lambda} e^{(\nu+\lambda)/2}\Big), \qquad (30)\end{aligned}$$

and then

$$\begin{aligned} H^1_1 &= -e^\lambda H^{11} \qquad (31)\\ &= -2\nu'((e^\lambda - 1)/r)(e^{-\lambda}/r)^2 - (e^\lambda - 1)^2(e^{-\lambda}/r^2)^2. \qquad (32)\end{aligned}$$

These expressions, Eq. (27) and Eq. (31), have been derived by a different method in 1985[3],[5], and were used for solving the Einstein equation. This process will be recapitulated in the next section. The difference from 1985 is that this time we have shown that this is the exact solution for a SSS metric.

III. QUANTUM CORRECTIONS FOR A SSS METRIC

The Einstein equation for a SSS metric, Eq. (7), is expressed

$$G^\mu_\rho = R^\mu_\rho - \delta^\mu_\rho R = -\frac{l^2\beta}{2} H^\mu_\rho, \qquad (33)$$

where[1]

$$l^2 = 16\pi G \qquad (34)$$

and

$$\beta = \frac{1}{2880\pi^2}(N_s + 11N_\nu + 62N_V), \qquad (35)$$

N_s, N_ν and N_V being the numbers of scalar fields, four-component neutrino fields and vector fields respectively. An explicit form of the Einstein equation reads

$$\begin{aligned} &e^{-\lambda}\big(\frac{\lambda'}{r} - \frac{1}{r^2}\big) + \frac{1}{r^2}\\ &= \frac{\xi}{2}\Big(e^{-2\lambda}\big(\frac{2\lambda'}{r^3} + \frac{1}{r^4}\big) + 2e^{-\lambda}\big(-\frac{\lambda'}{r^3} - \frac{1}{r^4}\big) + \frac{1}{r^4}\Big) \qquad (36)\end{aligned}$$

and

$$\begin{aligned} &e^{-\lambda}\big(\frac{-\nu'}{r} - \frac{1}{r^2}\big) + \frac{1}{r^2}\\ &= \frac{\xi}{2}\Big(e^{-2\lambda}\big(\frac{-2\nu'}{r^3} + \frac{1}{r^4}\big) + 2e^{-\lambda}\big(\frac{\nu'}{r^3} - \frac{1}{r^4}\big) + \frac{1}{r^4}\Big), \qquad (37)\end{aligned}$$

where

$$\xi = l^2\beta. \qquad (38)$$

Subtraction of the two Einstein equations yields

$$\nu' + \lambda' = 0 \qquad (39)$$

or

$$\nu + \lambda = 0 \qquad (40)$$

by the boundary condition at

$$r = \infty. \qquad (41)$$

Multiplying Eq. (36) by r^2,

$$\begin{aligned} e^{-\lambda}(r\lambda' - 1) + 1 = \frac{\xi}{2}\Big(&e^{-2\lambda}\big(\frac{2\lambda'}{r} + \frac{1}{r^2}\big)\\ &+ 2e^{-\lambda}\big(-\frac{\lambda'}{r} - \frac{1}{r^2}\big) + \frac{1}{r^2}\Big), \qquad (42)\end{aligned}$$

which can be expressed as

$$\begin{aligned} -(re^{-\lambda})' + r' = \frac{\xi}{2}\Big(&-\big(\frac{e^{-2\lambda}}{r}\big)'\\ &+ 2\big(\frac{e^{-\lambda}}{r}\big)' - \big(\frac{1}{r}\big)'\Big), \qquad (43)\end{aligned}$$

which can be integrated

$$\begin{aligned} &- re^{-\lambda} + r\\ &= \frac{\xi}{2}\big(-\frac{e^{-2\lambda}}{r} + 2\frac{e^{-\lambda}}{r} - \frac{1}{r}\big) + K, \qquad (44)\end{aligned}$$

or

$$e^{-2\lambda} - 2\big(1 + \frac{r^2}{\xi}\big)e^{-\lambda} + 1 + \frac{2r^2}{\xi} - \frac{2Kr}{\xi} = 0, \qquad (45)$$

where K stands for an integration constant. The solution of this quadratic equation is

$$e^{-\lambda} = e^\nu = 1 + \frac{r^2}{\xi} - \sqrt{\frac{r^4}{\xi^2} + \frac{2Kr}{\xi}}. \qquad (46)$$

This is the solution[3],[5] that was obtained by the author in 1985. In the limit, $r \to \infty$ or $\xi \to 0$, one obtains

$$e^{\nu} \to 1 - \frac{K}{r} + \frac{K^2\xi}{2r^4} + \cdots, \tag{47}$$

so that the integration constant K is determined to be

$$K = 2GM. \tag{48}$$

The important point is that this is the exact solution for quantum corrections for a SSS metric. As $r \to 0$,

$$e^{\nu} \to 1 - \sqrt{\frac{2Kr}{\xi}} \tag{49}$$

which shows the repulsive effect of quantum corrections on gravity at short distances. As is well known by now this is the basis for the author's model of high energy cosmic ray emission from AGN.

IV. APPENDIX: CURVATURE IN A SSS METRIC IN N DIMENSIONS

The extension of the curvature in a SSS metric to n dimensions reads

$$R^{01}_{....01} = (\frac{\nu''}{2} + \frac{\nu'^2}{4} - \frac{\nu'\lambda'}{4})e^{-\lambda} \tag{50}$$

$$\begin{aligned} R^{02}_{....02} &= R^{03}_{....03} = \cdots \\ &= R^{0,n-1}_{....0,n-1} = \frac{\nu'}{2}\frac{e^{-\lambda}}{r} \end{aligned} \tag{51}$$

$$\begin{aligned} R^{12}_{....12} &= R^{13}_{....13} = \cdots \\ &= R^{1,n-1}_{....1,n-1} = -\frac{\lambda'}{2}\frac{e^{-\lambda}}{r} \end{aligned} \tag{52}$$

$$\begin{aligned} R^{23}_{....23} &= R^{24}_{....24} = \cdots = R^{2,n-1}_{....2,n-1} = R^{34}_{....34} \\ &= \cdots = R^{n-2,n-1}_{....n-2,n-1} = -(e^{\lambda} - 1)\frac{e^{-\lambda}}{r^2} \end{aligned} \tag{53}$$

$$\begin{aligned} R^{0}_{..0} &= R^{01}_{....01} + R^{02}_{....02} + \cdots + R^{0,n-1}_{....0,n-1} \\ &= (\frac{\nu''}{2} + \frac{\nu'^2}{4} - \frac{\nu'\lambda'}{4})e^{-\lambda} + (n-2)\frac{\nu'}{2}\frac{e^{-\lambda}}{r} \end{aligned} \tag{54}$$

$$\begin{aligned} R^{1}_{..1} &= R^{10}_{....10} + R^{12}_{....12} + R^{13}_{....13} + \cdots + R^{1,n-1}_{....1,n-1} \\ &= (\frac{\nu''}{2} + \frac{\nu'^2}{4} - \frac{\nu'\lambda'}{4})e^{-\lambda} - (n-2)\frac{\lambda'}{2}\frac{e^{-\lambda}}{r} \end{aligned} \tag{55}$$

$$\begin{aligned} R^{2}_{..2} &= R^{20}_{....20} + R^{21}_{....21} + R^{23}_{....23} + \cdots + R^{2,n-1}_{....2,n-1} \\ &= \frac{\nu' - \lambda'}{2}\frac{e^{-\lambda}}{r} - (n-3)(e^{\lambda} - 1)\frac{e^{-\lambda}}{r^2} \\ &= R^{3}_{..3} = R^{4}_{..4} = \cdots = R^{n-1}_{..n-1} \end{aligned} \tag{56}$$

$$\begin{aligned} R &= R^{0}_{..0} + R^{1}_{..1} + \cdots + R^{n-1}_{..n-1} \\ &= (\nu'' + \frac{\nu'^2}{2} - \frac{\nu'\lambda'}{2})e^{-\lambda} + (n-2)(\nu' - \lambda')\frac{e^{-\lambda}}{r} \\ &- (n-2)(n-3)(e^{\lambda} - 1)\frac{e^{-\lambda}}{r^2} \end{aligned} \tag{57}$$

For the quadratic curvature form,

$$\begin{aligned} & R^{\alpha\beta}_{....\gamma\delta}R^{....\gamma\delta}_{\alpha\beta} \\ & = 4((R^{01}_{....01})^2 + (R^{02}_{....02})^2 + \cdots + (R^{0,n-1}_{....0,n-1})^2 \\ & + (R^{12}_{....12})^2 + (R^{13}_{....13})^2 + \cdots + (R^{1,n-1}_{....1,n-1})^2 \\ & + (R^{23}_{....23})^2 + (R^{24}_{....24})^2 + \cdots + (R^{2,n-1}_{....2,n-1})^2 \\ & + (R^{34}_{....34})^2 + \cdots \\ & + \cdots \\ & + (R^{n-2,n-1}_{....n-2,n-1})^2) \\ & = ((\nu'' + \frac{\nu'^2}{2} - \frac{\nu'\lambda'}{2})e^{-\lambda})^2 + (n-2)(\nu'^2 + \lambda'^2)(\frac{e^{-\lambda}}{r})^2 \\ & + 2(n-2)(n-3)((e^{\lambda} - 1)\frac{e^{-\lambda}}{r^2})^2 \end{aligned} \tag{58}$$

$$\begin{aligned} R^{\alpha}_{..\beta}R^{..\beta}_{\alpha} &= (R^{0}_{..0})^2 + (R^{1}_{..1})^2 + \cdots (R^{n-1}_{..n-1})^2 \\ &= \frac{1}{2}((\nu'' + \frac{\nu'^2}{2} - \frac{\nu'\lambda'}{2})e^{-\lambda})^2 \\ &+ \frac{n-2}{2}(\nu'' + \frac{\nu'^2}{2} - \frac{\nu'\lambda'}{2})(\nu' - \lambda')\frac{e^{-2\lambda}}{r} \\ &+ \frac{(n-2)^2}{4}(\nu'^2 + \lambda'^2)(\frac{e^{-\lambda}}{r})^2 \\ &+ \frac{(n-2)}{4}(\nu' - \lambda')^2(\frac{e^{-\lambda}}{r})^2 \\ &- (n-2)(n-3)(\nu' - \lambda')\frac{e^{\lambda} - 1}{r}(\frac{e^{-\lambda}}{r})^2 \\ &+ (n-2)(n-3)^2(\frac{e^{\lambda} - 1}{r})^2(\frac{e^{-\lambda}}{r})^2 \end{aligned} \tag{59}$$

and

$$\begin{aligned} R^2 &= ((\nu'' + \frac{\nu'^2}{2} - \frac{\nu'\lambda'}{2})e^{-\lambda} + (n-2)(\nu' - \lambda')\frac{e^{-\lambda}}{r} \\ &- (n-2)(n-3)(e^{\lambda} - 1)\frac{e^{-\lambda}}{r^2})^2. \end{aligned} \tag{60}$$

Acknowledgments

It is a great pleasure to thank David N. Williams for reading the manuscript.

---

[1] Birrell, N. D. and Davies, P. C. W., Quantum Fields in Curved Space (Cambridge University Press 1982).

[2] Anderson, P., Phys. Rev. D 28, 271 (1983); ibid., D 29, 615 (1984).

[3] Tomozawa, Y., Magnetic Monopoles, Cosmic Rays and Quantum Gravity, in the Proc. of 1985 INS International Symposium on Composite Models of Quarks and Leptons (Tokyo, edit. Terazawa, H. and Yasue, M., 1985), pp. 386.

[4] Tomozawa, Y., The Origins of Cosmic Rays and Quantum Effects of Gravity, in Quantum Field Theory (ed. Mancini, F., Ersever Science Publishers B. V., 1986) pp. 241. This book is the Proceedings of the International Symposium in honor of Hiroomi Umezawa held in Positano, Salerno, Italy, June 5-7, 1985.

[5] Tomozawa, Y., Cosmic Rays, Quantum Effects on Gravity and Gravitational Collapse, Lectures given at the Second Workshop on Fundamental Physics, University of Puerto Rico, Humacao, March 24-28, 1986. The lecture notes can be retrieved from KEK Kiss NO 200035789 at http://www-lib.kek.jp/KISS/kiss_prepri.html

[6] The Pierre Auger Collaboration, Science 318, 938 (2007); Correlation of the Highest-energy Cosmic Rays with the Positions of Nearby Active Galactic Nuclei, arXiv: 0712.2843 (2007).

[7] Tomozawa, Y., Evidence for a Dark Matter Particle, arXiv, astro-ph. 1002.1938 (2010).

[8] Tomozawa, Y., Cosmic Rays from AGN, the Knee Energy Mass Scale and Dark Matter Particles, arXiv, astro-ph. 1002.1327 (2010).

[9] Aharonian, F. et.al., Astron. & Astroph. 477, 353 (2008).